

## 衛生福利部臺東醫院高警訊藥品一覽表(108年9月)

1. 目的：利用各項提示功能，減少高警訊藥品發生用藥錯誤，以確保病人用藥安全。
2. 定義：高警訊藥品 (High Alert Medications)：當藥品使用錯誤時，有很高的危險會對病患造成明顯的傷害，即為高警訊藥品。
3. 依據2014年美國ISMP (Institute of Safety Medication Practices)公告、沒有固定規則化使用劑量、安全指數狹窄之藥品。
4. 本院高警訊藥品儲位與其他藥物區隔並以”紅色標籤”明顯標示，讓藥師於調劑時，有特別提醒注意之效果。
5. 高警訊藥品於藥袋上以”●”警示，以提醒藥師謹慎調劑並充分衛教病人。
6. 於針劑藥品上以”紅色”圓形標籤警示，以提醒醫療人員注意調劑及給藥。
7. 需冷藏之高警訊藥品應與其他藥品分層擺放。
8. 本院高警訊藥品有14類 (44項藥品)

<b>藥品分類</b>		
<b>Adrenergic agonists, IV</b>		
藥品學名	商品名	規格
Epinephrine	Adrenalin	1 mg/ml/amp.
Dopamine	Gipamine	600 mg/200 ml/bag
Norepinephrine	Norepinephrine	4mg/4ml/amp.
<b>Adrenergic antagonists, IV</b>		
Labetalol	Trandate	25 mg/5 ml/amp.
<b>Antiarrhythmics, IV</b>		
Amiodarone	Cardarone	150 mg/3 ml/amp.
Adarone(口服)	Amiorone	200 mg/tab.
Adenosine	Adenocor	6 mg/2 ml/vial
Lidocaine	2% Xylocaine	2%, 20 ml/vial
Atropine	Atropine	1 mg/ml/amp.
<b>Antithrombotic agents, including: Anticoagulants</b>		
Warfarin	Orfarin	5 mg/tab.
Heparin	Agglutex	25000 IU/5 ml/vail
<b>Dextrose, hypertonic, 20% or greater</b>		
50% Glucose	Vitagen	50%, 20 ml/amp.
<b>Inotropic medications, IV</b>		
Digoxin	Lanoxin	0.5 mg/2 ml/amp.
Digoxin (口服)	Lanoxin	0.25 mg/tab.
<b>Hypoglycemics, oral</b>		
Glimepiride	Amalin	4 mg/tab.

Repaglinide	Repaglinide	1 mg/tab.
Metformin	C.T. L XR	500 mg/tab.
Metformin	Henformin	850 mg/tab.
Pioglitazone	Glitis	30 mg/tab.
Sitagliptin	Januvia	100 mg/tab.
Alogliptin	NESINA	25 mg/tab.
Miglitol	Migbose	50 mg/tab.
Pioglitazone- Metformin	Lodigit	15-850 mg/tab.
Glimepiride-Metformin	Glimet	2-500 mg/tab.
<b>Insulin, subcutaneous and IV</b>		
Insulin, Regular	Humulin R	1000 IU/10 ml/vial
Insulin aspart	NovoRapid	300 IU/3 ml /FlexPen
30% Soluble insulin aspart-70% Insulin aspart crystallised with protamine	NovoMix 30	300 IU/3 ml/FlexPen
Insulin Detemir	Levemir	300 IU/3 ml/FlexPen
<b>Moderate sedation agents, IV</b>		
Midazolam	Midatin	15 mg/3 ml/amp.
<b>Sedative &amp; Hypnotic injection</b>		
Diazepam	Dianlin	10 mg/2 ml/amp.
Lorazepam	Anxicam	2 mg/ml/amp.
Pnenytoin	Aleviatin	250 mg/5 ml/amp.
Pnenytoin (口服)	Aleviatin	100 mg/tab.
<b>Chemotherapeutic agnet</b>		
Methotrexate	Methotrexate	2.5 mg/tab.
<b>Narcotics/opioids, IV and oral</b>		
Pethidine	Pethidine	50 mg/ml/amp.
<b>Sodium chloride for injection, hypertonic, greater than 0.9% concentration</b>		
3% Sodium chloride	3% Sodium chloride	15 gm/500 ml/bot.
<b>Special medications</b>		
Nitroglycerin sublingual tab.	Nitrostat sublingual tab.	0.6mg/ sublingual tab.
Nitroglycerin	Millisrol inj.	5 mg/10 ml/amp.
Magnesium sulfate	10% Magnesium sulfate	16.23 mEq/20 ml/amp.
Potassium Chloride- Riboflavin	K.C.L	3 gm(40mEq)- 6 mg/20 ml/amp.

Sodium bicarbonate	Rolikan	7%, 20 ml/amp. ( HCO <sub>3</sub> <sup>-</sup> 16.67mEq/20 ml)
Aminophyllin	Aminophyllin	250 mg/10 ml/amp.
Aminophylline	Aminophylline	100 mg/tab.
Calcium gluconate	2% Vitacal	Ca <sup>2+</sup> 5.44mg/ml,108.8mg/ 20 ml/amp.

資料來源：1. [ISMP](#) List of **High-Alert** Medications in Acute Care Settings ISMP 2014.

([www.ismp.org](http://www.ismp.org))

2. TSHP 高警訊藥品清單(草案) 2018.04.18